

A singular perturbation approach to vector-borne epidemic models

Fred Brauer¹

¹ *Department of Mathematics, University of British Columbia, Vancouver, BC V6T 1Z2, Canada* brauer@math.ubc.ca

² *University of Florida, Gainesville, FL, US, 32611*

In vector-borne epidemic models there is often a substantial difference between the vector and host time scales. This makes it possible to use the quasi-steady-state to obtain final size relations.

References

- [1] Brauer, F. (2017)
A final size relation for epidemic models of vector - transmitted diseases, *Infectious Disease Modelling*, **2**: 12–20.
- [2] Hoppensteadt, F.C. (1966)
Singular perturbations on the infinite interval, *Trans. Amer. Math. Soc.* **123**: 521–535.
- [3] Levinson, N. (1950)
Perturbations of discontinuous solutions of nonlinear systems of differential equations, *Acta Math.* **82**: 71–106.
- [4] Rocha, F., M. Aguiar, M. Souza, & N. Stollenwerk (2013)
Time-scale separation and centre manifold analysis describing vector-borne disease dynamics, *Int. J. Comp. Math* **90**: 2105–2125.
- [5] Segel, L.A. & M. Slemrod (1989)
The quasi-steady-state assumption: A case study in perturbation, *SIAM Review* **31**: 446–477.

*Mini-Symposium: Mathematical Epidemiology

- [6] Tihonov, A.N. (1948)
On the dependence of the solutions of differential equations on a small parameter, *Mat. Sbornik NS* **22**; 193–204.
- [7] van den Driessche, P. and J. Watmough (2002)
Reproduction numbers and subthreshold endemic equilibria for compartmental models of disease transmission, *Math. Biosc.* **180**:29–48.