

Ebola: Impact of hospitals admission policy in an overwhelmed scenario

Mondal Hasan Zahid and Cristopher Kribs

University of Texas at Arlington, 411 S Nedderman Drive, Arlington, TX, US,
76019-0408 mondalhasan.zahid@mavs.uta.edu
kribs@uta.edu

Infectious disease outbreaks sometimes overwhelm healthcare facilities. A recent case occurred in West Africa in 2014 when an Ebola virus outbreak overwhelmed facilities in Sierra Leone, Guinea and Liberia. In such scenarios, how many patients can hospitals admit to minimize disease burden? This study considers what type of hospital admission policy during a hypothetical Ebola outbreak can better serve the community, if overcrowding degrades the hospital setting. Our result shows that which policy minimizes loss to the community depends on the initial estimation of the control reproduction number, R_0 . When the outbreak grows extremely fast ($R_0 \gg 1$) it is better (in terms of total disease burden) to stop admitting patients after reaching the carrying capacity because overcrowding in the hospital makes the hospital setting ineffective at containing infection, but when the outbreak grows only a little faster than the system's ability to contain it ($R_0 \gtrsim 1$), it is better to admit patients beyond the carrying capacity because limited overcrowding still reduces infection more in the community. However, when R_0 is no more than a little greater than 1 (for our parameter values, 1.012), both policies result the same because the number of patients never exceeds the maximum capacity.

References

- [1] J. M. Drake et. al, Ebola cases and health system demand in liberia, *PLoS Biology*, **13** (2015), e1002056.
- [2] WHO, *Health Statistics and Information Systems*, Update date: 07-01-2016, Access date: 03-03-2017.
- [3] WHO, *Liberia: Ebola Treatment Centre Sets A New Pace*, October 2014, Access date 2018-06-01.

*Mini-Symposium: Dynamics of infectious diseases and public health